
TummyCare-Max


Simple Mixing Guide

For use with generic omeprazole OTC tablets 20mg at a concentration of 4 mg / mL concentration

Your kit includes:



Step 2: Add water to bottle




Step 1: Add Prilosec® to grinder

Put 28 generic omeprazole OTC 20mg tablets into the coffee grinder with 1 tspn TCMax powder, grind until pulverized (~ 8 pulses 10 seconds each, shake the grinder up and down while holding lid tightly to ensure a fine grind of tablets.
Using the funnel, pour into the bottle, using a small spoon or spatula to get all powder from coffee grinder Using funnel, pour the rest of the powder from the TummyCare Max packet into the bottle.

Coffee grinder such as Krups, Toastmaster or Quiseen (see below). The Quissen is rated very high and is \$9.99



Step 3: Dispense



Remember:

- Refrigerate between doses.
- Do not give with acidic foods (fruit, fruit juice, fruit yogurt).
- Use within 28 days of preparing the suspension.
- The kits are reusable. Please save the bottle, syringes, and funnel for future use. You will need only to reorder refill packets.

Recommended Dosing

Total volume	4 mg/mL
dispensed	concentration
4.0mL	16 mg
4.5mL	18 mg
5.0mL	20 mg
5.5mL	22 mg
6.0mL	24 mg
6.5mL	26 mg
7.0mL	28 mg
7.5mL	30 mg
8.0mL	32 mg
8.5mL	34 mg
9.0mL	36 mg

- 1mL = 1cc
- 3 mL is the minimum volume per dose of TummyCare Max that should be given. (This is to ensure enough buffer to neutralize stomach acid)
- Doses can be given regardless of meal-time.
- For more information on dosing, visit www.infant-acid-reflux-solutions.com.