

TummyCare-max


Simple Mixing Guide

For use with Prevacid® 30 mg Capsules
at a concentration of 3 mg/mL

Your kit includes:


1. capped bottle
2. dosing syringe
3. filling syringe
4. funnel

Kits are for reuse. Do NOT throw away caps, bottles, funnel, syringes as you will reuse these.



Step 1: Add Prevacid® granules

- Set aside 15 Prevacid 30 mg capsules
- Moisten tips of capsules with wet cloth
- Holding the capsule over the funnel, snip the top of the capsule with scissors; empty the contents of each of the 15 capsule of 30mg Prevacid. into the grinder.



Step 2: Add 1 tspn TC Max to grinder


Grind until pulverized (~ 8 pulses 10 seconds each, shake the grinder up and down while holding lid tightly to ensure a fine grind of tablets.

- Using the funnel, pour into the bottle, using a small spoon or spatula to get all powder from coffee grinder
- Using funnel, pour the rest of the powder from the TummyCare Max packet into the bottle.




Step 3: Add water to bottle

- Using syringe, add 108 mL (60mL, then 48mL) tap water to bottle -- do NOT use liquids with provided funnel
- Immediately cap and shake bottle
- Refrigerate 15min before step 3



Step 4: Dispense

- Shake bottle before each dose
- Secure the dosing syringe to the blue cap
- Turn the bottle up-side-down and draw the dose volume into the dosing syringe
- Administer to your child



Remember:

- Refrigerate between doses.
- Do not give with acidic foods (fruit, fruit juice, fruit yogurt).
- Use within 28 days of preparing the suspension.
- The kits are reusable. Please save the bottle, syringes, and funnel for future use. You will need only to reorder refill packets.

Coffee grinder such as Krups, Toastmaster or Quiseen (see below). The Quiseen is rated very high and is \$9.99



Recommended Dosing

Total volume dispensed	3mg/mL concentration
4.0mL	12mg
4.5mL	13.5mg
5.0mL	15mg
5.5mL	16.5mg
6.0mL	18mg
6.5mL	19.5mg
7.0mL	21mg
7.5mL	22.5mg
8.0mL	24mg
8.5mL	25.5mg
9.0mL	27mg

- 1mL = 1cc
- 3 milliliters (mL) is the minimum volume per dose of the TummyCare-Max™ that should be given.
- Doses can be given regardless of meal-time.
- For more information on dosing, visit www.infant-acid-reflux-solutions.com